arXiv:1410.0399v2 [quant-ph] 2 Mar 2016

2D Schrödinger Equation with Mixed Potential in Noncommutaive Complex space

Slimane Zaim and Hakim Guelmamene
Département de Physique ,
Faculté des Sciences de la Matiére, Université Batna1, Algeria.

Abstract

We study the 2D Schrdinger equation for Hydrogen atom with the lenear plus Harmonic Potentials in noncommutative complex space, using the Power-series expansion method and perturbation theory. Hence we can say that the Schrödinger equation in noncommutative complex space describes to the particles with spin (1/2) in an external uniform magnitic field. Where the noncommutativity play the role of magnetic field with created the total magnetic moment of particle with spin 1/2, who in turn shifted the spectrum of energy. Such effects are similar to the Zeeman splitting in a commutative space. It is shown that on the noncommutative space the degeneracy of the levels is lifted completely.

Keywords:Central potential, Noncommutative Geometry, complex space,
Pacs numbers:11.10.Nx, 32.30-r, 03.65-w

1 introduction:

The non-commutative quantum mechanics is motivated by the natural extension of the commutation relations between position and momentum, by imposing further commutation relations between position coordinates themselves. the noncommutativity of position coordinates immediately implies a set of uncertainty relations between position coordinates analogous to the Heisenberg uncertainty relations between position and momentum; namely:

$$\Delta\hat{x}_i\Delta\hat{x}_j \succeq \frac{\theta_{ij}}{2} \tag{1}$$

Using the new noncommutative coordinate operators $\hat{x}$ satisfying the commutation relations:

$$[\hat{x}_i, \hat{x}_j] == i\theta_{ij} \tag{2}$$

Where $\hat{x}^i = x^i - \frac{\theta^{ij}}{2}p_j$ and the parameter θ^{ij} is an antisymmetric real matrix of dimension square length in the noncommutative canonical-type space. Where

the ordinare product betwenn functions in noncommutive space is remplaced by the star product, where the star product between two arbitrary functions $f(x)$ and $g(x)$ is given by:

$$f(x) \star g(x) = f(x).g(x) + \exp\left(\frac{i}{2}\theta^{ij}\partial_i\acute{\partial}_j\right) f(x)\, g(\acute{x})\Big|_{x=\acute{x}} \tag{3}$$

We observed that the result of eq. (1) satisfied by applying the notion of the star product represented by eq.(2). The important problems of noncommutative quantum mechanics is to solve the Schrodiger equation with physical potentials in noncommutative space. Actually there are many attempts to study noncommutative space but in two dimensional complex space the study is a limited attempts [1, 2]. The study of the Schrödinger equation for Hydrogen atom with a linear plus Harmonic potentials has proved to be fruitful and many papers have been published $[3-12]$. In this work we study an important contribution to the non-commutative geometry approach to the hydrogen atom with linear plus Harmonic potentials. Our goal is to solve the Schrödinger equation for Hydrogen atom with linear plus Harmonic potentials in a non-commutative complex space up to first-order of the non-commutativity parameter using the Moyal product , we show that the non-commutative value of the energy levels indicates the splitting of $1s$ states.

This paper is organized as follows. In section 2 we derive the deformed $2D$ Schrödinger equation for Hydrogen atom with a linear plus Harmonic potentials in noncommutative space, we solve this equation and obtain the noncommutative modification of the energy levels. In section 3 we derive the deformed $2D$ Schrödinger equation for Hydrogen atom with a linear plus Harmonic potentials in noncommutative complex space, we solve this equation and obtain the non-commutative modification of the energy levels. Finally, in section 4, we draw our conclusions.

2 Noncommutative Schrödingre equartion with mixed potential

The $2D$ Schrödinger equation with the central potential $V(\hat{r})$, in noncommutative space, which have the following form:

$$\left(-\frac{1}{2m}\Delta + V(r)\right) * \Psi(r,\varphi) = \hat{E}\Psi(r,\varphi) \tag{4}$$

Expanding the $*$ product for equ.(3) we get

$$\left(-\frac{1}{2m}\Delta + V(\hat{r})\right)\Psi(r,\varphi) = \hat{E}\Psi(r,\varphi)$$

where the $V(\hat{r})$ (coulomb plus linear plus oscilator harmonic) in nonocommutative space as :

$$V(\hat{r}) = \frac{c}{\hat{r}} + a\hat{r} + b\hat{r}^2 \tag{5}$$

and the deformed of $\hat{r}$ to first order in the nonocommutative parameter θ as:

$$\hat{r} = r - \frac{\theta \cdot L}{2r} + \mathcal{O}\left(\Theta^2\right) \tag{6}$$

Then the deformed of the central potential $V(\hat{r})$ in eq.(5) up to $\mathcal{O}\left(\Theta^2\right)$ as:

$$\begin{aligned} V(\hat{r}) &= ar + br^2 + \frac{c}{r} - \left(\frac{a}{2} + b\right)\theta \cdot L + \frac{c\theta \cdot L}{2r^3} + \mathcal{O}\left(\Theta^2\right) \\ &= V(r) + V_{\mathrm{NC}}^{\theta}(r) + \mathcal{O}\left(\Theta^2\right) \end{aligned} \tag{7}$$

where

$$V_{\mathrm{NC}}^{\theta} = -\left(\frac{a}{2} + b\right)\theta \cdot L + \frac{c\theta \cdot L}{2r^3} \tag{8}$$

wthere the first term (not depeneds on the distance r), which is similar to the interaction between the spin and angular momentum, where the non-commutativity leads the role of the spin and the second term is represent the dipole-dipole interaction and is the leading-order perturbation . For simplicity, first of all, we choose the coordinate system (x, y, z) so that $\theta^{xy} = -\theta^{yx} = \theta\delta^{xy} = \theta_z$, such that $\theta \cdot L = \theta L_z$ and assume that the other components are all zero ,then the noncommutative Hamiltonian up to $\mathcal{O}\left(\Theta^2\right)$ as the form:

$$\hat{H} = \hat{H}_O + H_{\mathrm{pert}}^{\theta} \tag{9}$$

where

$$\hat{H}_O = H_O - \theta\left(\frac{a}{2} + b\right)\theta L_z \text{ and } \; H_{\mathrm{pert}}^{\theta} = \frac{c\theta L_z}{2r^3} \tag{10}$$

To investigate the modification of the energy levels by eq. (10), we use the first-order perturbation theory. The spectrum of $\hat{H}_0$ and the corresponding wave functions are well-known and given by [3]:

$$\begin{cases} \Phi(\varphi) = \exp(\pm im\varphi) \text{ where } m = 0; 1, 2... \\ R_m^n(r) = \exp\left(-\frac{ar+br^2}{2\sqrt{b}}\right)\sum_{p=0}^{n} a_p r^{p+\delta} \end{cases}, \tag{11}$$

and

$$\hat{E}_{n,m} = 2\sqrt{b}(1 + m + n) - \frac{\theta}{2}(a + 2b)m \tag{12}$$

Now, the correction to the energy to first order in θ is:

$$\begin{aligned}\Delta E_{n,m}^{\theta} &= \langle n|\, H_{\text{pert}}^{\theta}(r)\,|n\rangle = \theta m \int \Psi^{(n)*}(r)\,\frac{c}{2r^3}\Psi^{(n)}(r)\,r dr d\varphi \\ &= c\pi\theta m \int R^{(n)*}(r)\,\frac{1}{r^2}R^{(n)}(r)\,dr \end{aligned} \tag{13}$$

where:

$$\begin{aligned}\int R^{(n)*}(r)\,\frac{1}{r^2}R^{(n)}(r)\,dr &= \sum_{p=0}^{n} C_p \int r^{p+2\delta-2}\exp\left(-\frac{ar+br^2}{2\sqrt{b}}\right)dr \\ &= \exp\left(\frac{a^2+2ba}{4b\sqrt{b}}\right)\sum_{p=0}^{n} C_p b^{-\frac{n+2\delta}{2}}\Gamma\left(n+2\delta,-\frac{a}{2\sqrt{b}}\right)\end{aligned} \tag{14}$$

where

$$C_p = \sum_{k=0}^{p} a_k a_{p-k} \tag{15}$$

Then the modification of the energy levels is given by:

$$\Delta E_{n,m}^{\theta} = \theta\left(c\pi\exp\left(\frac{a^2+2ba}{4b\sqrt{b}}\right)\sum_{p=0}^{n} C_p b^{-\frac{n+2\delta}{2}}\Gamma\left(n+2\delta,-\frac{a}{2\sqrt{b}}\right)\right)m \tag{16}$$

We have show that the non-commutative energy correction is proportional to the quantum number m , such are similar to the Zeeman effects and they remove the degeneracy with respect to m.

3 Schrödingre equartion with mixed potential in noncommutative complex space

In the complex coordinate $(z,\bar{z})$, the radial wave function $\Psi(r,\varphi)$ will be written to the form:

$$\Psi(r,\varphi) = \Psi(z,\bar{z}) = \frac{R_m(z,\bar{z})}{(z\bar{z})^{1/4}}\Phi(\varphi), \qquad z\bar{z} = r^2 \tag{17}$$

where the energy levels and the corresponding radial wave functions are well-known and given by [3, 8]:

$$.R_m(z,\bar{z}) = \exp\left(p_m(z,\bar{z})\right)\sum_{p=0}^{n} a_n\left(z\bar{z}\right)^{\frac{p+\delta}{2}} \tag{18}$$

and

$$E_n = 2\sqrt{b}\,(1+m+n) \tag{19}$$

where $p_m\,(z,\bar{z})$ is given by:

$$p_m\,(z,\bar{z}) = \alpha\sqrt{z\bar{z}} + \frac{1}{2}\beta z\bar{z} \tag{20}$$

and α , β and δ are given by :

$$\left\{ \begin{array}{l} \beta^2 = b, \quad 2\alpha\beta = a, \end{array} \right. \tag{21}$$

if we choose $\beta = -\sqrt{b}$ and $\delta = \frac{1}{2} + m$ .The exact solution for $n = 0$ and $n = 1$ are given by [3, 8]:

$$\left\{ \begin{array}{ccc} R_m^{(0)} = a_0\,(z\bar{z})^{\delta/2} \exp\left(-\frac{a(z\bar{z})^{1/2}+bz\bar{z}}{2\sqrt{b}}\right) & , & E_{0,m} = 2\sqrt{b}\,(1+m) \\ R_m^{(1)} = \left(a_0 + a_1\sqrt{z\bar{z}}\right)(z\bar{z})^{\delta/2} \exp\left(-\frac{a(z\bar{z})^{1/2}+bz\bar{z}}{2\sqrt{b}}\right) & , & E_{1,m} = 2\sqrt{a}\,(2+m) \end{array} \right. \tag{22}$$

In the noncommutative complex space we can show that, in the first order of the parameter θ:

$$\begin{array}{rcl} \hat{z}\hat{\bar{z}} & = & z\bar{z} - \theta\,(L_z - 1) \\ \hat{\bar{z}}\hat{z} & = & z\bar{z} - \theta\,(L_z + 1) \\ \hat{p}^2 & = & 4p_z p_{\bar{z}} = p^2 \end{array} \tag{23}$$

Then the noncommutative Hamiltonian $\hat{H}$ associated by the mixed potential as given by[2]:

$$\hat{H} = \frac{2}{m} p_{\hat{z}} p_{\hat{\bar{z}}} + V\left(\hat{z},\hat{\bar{z}}\right) \tag{24}$$

where the deformed potential $V\left(\hat{z},\hat{\bar{z}}\right)$ in the noncommutative complex coordinate $\left(\hat{z},\hat{\bar{z}}\right)$ as:

$$V\left(\hat{r}\right) = \left(\begin{array}{cc} V\left(\hat{z},\hat{\bar{z}}\right) = a\sqrt{\hat{z}\hat{\bar{z}}} + b\hat{z}\hat{\bar{z}} + \frac{c}{\sqrt{\hat{z}\hat{\bar{z}}}} & 0 \\ 0 & V\left(\hat{\bar{z}},\hat{z}\right) = a\sqrt{\hat{\bar{z}}\hat{z}} + b\hat{\bar{z}}\hat{z} + \frac{c}{\sqrt{\hat{\bar{z}}\hat{z}}} \end{array} \right) \tag{25}$$

Then $V\left(\hat{z},\hat{\bar{z}}\right)$ and $V\left(\hat{\bar{z}},\hat{z}\right)$ as follows:

$$V\left(\hat{z},\hat{\bar{z}}\right) = V\,(r) + V_{NC}^{+}\,(r) \tag{26}$$

$$V\left(\hat{\bar{z}},\hat{z}\right) = V\,(r) + V_{NC}^{-}\,(r) \tag{27}$$

where

$$\begin{aligned} V_{NC}^{\pm}(r) &= -\theta\left(\frac{a}{2}+b-\frac{c}{2r^3}\right)(L_z \mp 1) \\ &= -\theta\left(\frac{a}{2}+b\right)(L_z-2s_z)+\theta\frac{c\,(L_z-2s_z)}{2r^3}, s_z=\pm\frac{1}{2} \end{aligned} \tag{28}$$

The noncommutative Hamiltonian in eq.(24) as:

$$\hat{H}=H_O-\theta\left(\frac{a}{2}+b\right)(L_z-2s_z)+\theta\frac{c\,(L_z-2s_z)}{2r^3} \tag{29}$$

where the first term H_O is the ordinary Hamiltonian of hydrogen atom with linear plus Harmonic potentials , the second term that not depeneds on the distance r, wisch is simelar to the interaction between the spin and angular momentum and the therd term is represent the dopol-dipol iteraction and is the leading-order perturbation . Then the noncommutative Hamiltonian in eq.(29) we can rewiteen as the form:

$$\hat{H}=\hat{H}_O+H_{\text{pert}}^{\theta} \tag{30}$$

where

$$\hat{H}_O=H_O-\theta\left(\frac{a}{2}+b\right)(L_z-2s_z) \tag{31}$$

and

$$H_{\text{pert}}^{\theta}=\theta\frac{c\,(L_z-2s_z)}{2r^3} \tag{32}$$

To investigate the correction of the energy levels by eq. (32), we use the first-order perturbation theory. The spectrum of $\hat{H}_0$ and the corresponding eigenfunctions can be written as:

$$R_m^{+}(z,\bar{z}) = \exp\left(p_m(z,\bar{z})\right)\sum_{n=0}a_n\left(z\bar{z}\right)^{\frac{n}{2}+\delta}\oplus|+\succ \tag{32}$$

and

$$R_m^{-}(z,\bar{z}) = \exp\left(p_m(z,\bar{z})\right)\sum_{n=0}a_n\left(z\bar{z}\right)^{\frac{n}{2}+\delta}\oplus|-\succ \tag{33}$$

where the energy levels are given by:

$$\hat{E}_{n;m}^{\pm}=2\sqrt{b}\,(1+m+n)-\theta\left(\frac{a}{2}+b\right)(m\mp 1) \tag{34}$$

Now to obtain the correction energy levels as a result of the eq.(31), we use perturbation theory. To calculate $\Delta E_{n,m}^{\theta\pm}$ associate withe the spin up and spin down , we use the radial function in equations (32) and (33), to obtain:

$$\begin{aligned} \Delta E_{n,m}^{\theta\pm} &= \langle n|\,H_{\text{pert}}^{\theta}(r)\,|n\rangle=\frac{\theta}{2}(m\mp 1)\int\Psi^{(p)*}(z,\bar{z})\frac{c}{z\bar{z}\sqrt{z\bar{z}}}\Psi^{(p)}(z,\bar{z})\,rdr \\ &= \theta\pi(m\mp 1)\int R^{(p)*}(r)\frac{c}{r^3}R^{(p)}(r)\,rdr \end{aligned} \tag{35}$$

The non-commutative modification of the energy levels to the stationary state $\Delta E_{0,m}^{\theta\pm}$ to the first order in θ as given by:

$$\begin{aligned}\Delta E_{0,m}^{\theta\pm} &= \pi\left(m\mp 1\right)\theta\int_{0}^{+\infty}\left[a_0 r^{\delta}\exp\left(-\frac{ar+br^2}{2\sqrt{b}}\right)\right]^2\frac{c}{r^3}rdr \\ &= \theta\left(m\mp 1\right)C_0 \end{aligned} \tag{36}$$

where :

$$\begin{aligned}C_0 &= \pi c a_0^2\int_{0}^{+\infty} r^{2\delta-2}\exp\left(-\frac{ar+br^2}{\sqrt{b}}\right)dr \\ &= \exp\left(\frac{a^2+2ba}{4b\sqrt{b}}\right)a_0^2 b^{-\delta}\Gamma\left(2\delta,-\frac{a}{2\sqrt{b}}\right)\end{aligned} \tag{37}$$

The noncommutative energy levels to the stationary state as:

$$\hat{E}_{0,m}^{\pm} = 2\sqrt{b}\left(1+m\right)+\left(C_0-\frac{a}{2}-b\right)\theta\left(m\mp 1\right) \tag{38}$$

We show that the non-commutative value of the energy levels indicates the splitting of $1s$ states in two levels. Also we can say that the Lamb shift is actually induced by the space non-commutativity which plays the role of a magnetic field and spin in the same moment (Zemann effect).

Now, the noncommutative energy levels to the excitation state n, to first order in θ as:

$$\begin{aligned}\hat{E}_{n,m}^{\pm} &= 2\sqrt{b}\left(1+m+n\right)+\theta\left(a-\frac{b}{2}\right)\left(m\pm 1\right)-2\left(m\pm 1\right)\theta\times \\ &\quad\int_{0}^{+\infty}\left[\left(a_0+a_1r^2+...a_pr^{2p}\right)\right]^2\left[r^{\delta-1}\exp\left(-\frac{ar+br^2}{2\sqrt{b}}\right)\right]^2 dr \\ &= 2\sqrt{b}\left(1+m+n\right)+\theta\left(a-\frac{b}{2}-2A_n\right)\left(m\pm 1\right)\end{aligned} \tag{39}$$

where

$$A_n = \exp\left(\frac{a^2+2ba}{4b\sqrt{b}}\right)\sum_{k=0}^{n}C_p b^{-\frac{k+2\delta}{2}}\Gamma\left(k+2\delta,-\frac{a}{2\sqrt{b}}\right) \tag{40}$$

This result is very important, as a possible means of introducing electron spin we replace $l=\pm\left(j+1/2\right)$and $n\to n-j-1-1/2$, where j is the quantum number associated to the total angular momentum. Then the $l=0$ state has the same total quantum number $j=1/2$. In this case the non-commutative value of the energy levels indicates the splitting of $1s$ states.

4 Conclusion

In this paper we started from quantum Hydrogen atom with the linear plus Harmonic potentials in a canonical non-commutative complex space, we have derived the defermed Schrodinger equation. Using the power series expansion method to solving and we found that the noncommutative energy levels, the energy levels are shifted to $(2j+1)$ levels ,it acts here like a Lamb shift in Dirac theory. This proofs that the non-commutativity has an effect similar to the Zeeman effects which iduced by the magnetic field [13] , where the non-commutativity leads the role of the magnetic field.

References

[1] Arbab, A.I.: Europhys. Lett. 98, 30008 (2012)

[2] S.Zaim, The quantum and the Klein-Gordon oscillators in a non -commutative complexe space and the thermodynamic functions, Int. J. Theor. Phys,(2014),[doi:10.1007/s10773-014-2007-y].

[3] Shi-. Hai Dong, Exact solutions of the two -dimensional Schrödinger equation with certain potentials, International Journal of Theoretical Physics. Vol.39 ,No.4, (2000).

[4] R. J. LeRoy and W. Lam, Chem. Phys. Lett. 71, 544 (1970); R. J. LeRoy and R. B. Bernstein, J. Chem. Phys. 52, 3869 (1970).

[5] E. Vogt and G. H. Wannier, Phys. Rev. 95, 1190 (1954).

[6] L. D. Landau and E. M. Lifshitz, Quantum Mechanics, Vol. 3, 3rd Ed. (Pergamon

Press, Oxford, 1977); D. R. Bates and I. Esterman, Advances in Atomic and Molecular

Physics, Vol. 6 (Academic, New York, 1970).

[7] B. H. Bransden and C. J. Joachain, Physics of Atoms and Molecules (Longman, London, 1983).

[8] Shi-. Hai Dong and all, Exact solutions of the Schrödinger equation with Singular Even-Power potential in two dimenstions, Found.Phys.Lett. 12 (1999) 465-474

[9] L. Witten, Gravitation: An introduction to Current Research. Wiley, New York (1962).

[10] B. K. Cheng and M. G. da lus, J. Phys. A. 24,2033(1992).

[11] G. C. Maitland, M. Rigby, E. B. Smith and W. A. Wakeham, Intermolecular Forces (Oxford University Press, Oxford, 1987).

[12] S. Zaim, Exact solutions of the 2D Schrödinger equation with central potentials induced by the non-commutativity of space,arXiv:1405.1002.